Magnetization and energy dynamics in spin ladders: Evidence of diffusion in time, frequency, position, and momentum

Jonas Richter,^{1,*} Fengping Jin,² Lars Knipschild,¹ Jacek Herbrych,^{3,4} Hans De Raedt,⁵ Kristel Michielsen,^{2,6} Jochen Gemmer,¹ and Robin Steinigeweg^{1,†}

¹Department of Physics, University of Osnabrück, D-49069 Osnabrück, Germany

²Institute for Advanced Simulation, Jülich Supercomputing Centre, Forschungszentrum Jülich, D-52425 Jülich, Germany

³Department of Physics and Astronomy, The University of Tennessee, Knoxville, Tennessee 37996, USA

⁴Materials Science and Technology Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

⁵Zernike Institute for Advanced Materials, University of Groningen, NL-9747 AG Groningen, Netherlands

⁶RWTH Aachen University, D-52056 Aachen, Germany

(Received 14 November 2018; revised manuscript received 15 February 2019; published 23 April 2019)

The dynamics of magnetization and energy densities are studied in the two-leg spin-1/2 ladder. Using an efficient pure-state approach based on the concept of typicality, we calculate spatiotemporal correlation functions for large systems with up to 40 lattice sites. In addition, two subsequent Fourier transforms from real to momentum space as well as from the time to frequency domain yield the respective dynamical structure factors. Summarizing our main results, we unveil the existence of genuine diffusion for both spin and energy. In particular, this finding is based on four distinct signatures which can all be equally well detected: (i) Gaussian density profiles, (ii) time-independent diffusion coefficients, (iii) exponentially decaying density modes, and (iv) Lorentzian line shapes of the dynamical structure factor. The combination of (i)–(iv) provides a comprehensive picture of high-temperature dynamics in this archetypal nonintegrable quantum model.

DOI: 10.1103/PhysRevB.99.144422

I. INTRODUCTION

The study of low-dimensional spin systems is one of the most active fields in condensed-matter physics. On the one hand, quantum spin models are of immediate relevance to describe the properties of various Mott insulators, where (quasi-)one-dimensional structures like chains or ladders are realized within the bulk materials. The notion of property can be manifold in this context, including thermodynamic quantities [1], transport characteristics such as the heat conductivity [2,3], and other dynamic features probed by, e.g., neutron scattering [4,5], nuclear magnetic resonance [6], and muon spin resonance [7], to name just a few. Particularly, developing a thorough understanding of quantum magnets, both experimentally and theoretically, is also of essential importance in order to pave the way for potential spin-based technologies in the future [8].

On the other hand, from a more fundamental point of view, low-dimensional spin models represent prototypical examples of interacting quantum many-body systems, allowing us to study questions ranging from the foundations of statistical mechanics [9] to the physics of black holes [10]. In particular, long-standing questions concerning the emergence of thermodynamic behavior in isolated quantum systems have recently experienced rejuvenated attention [11–14]. This upsurge of interest is due to not least the advance of controlled experiments with cold atoms and trapped ions [15–17], theoretical key concepts such as the eigenstate thermalization hypothesis [18–21] and the typicality of pure quantum states [22–25], and the development of powerful numerical techniques [26].

Concerning the relaxation in isolated quantum systems, an intriguing question is the difference in the properties between integrable and nonintegrable systems. On the one hand, integrable systems exhibit a macroscopic set of (quasi)local conservation laws [27,28] which might cause anomalous thermalization [29] as well as nondecaying currents, i.e., ballistic transport [30]. Nonetheless, even for paradigmatic integrable models, signatures of diffusion have been reported, e.g., for the spin-1/2 XXZ chain above the isotropic point [31–36] and the Fermi-Hubbard model for strong particle-particle interactions [37-39]. Moreover, significant progress in understanding transport in integrable models was recently achieved within the framework of generalized hydrodynamics [40-42]. On the other hand, in more realistic situations, integrability is often lifted due to various perturbations, e.g., spin-phonon coupling [43], long-range interactions [44], dimerization [45], and the presence of impurities [46] or disorder [47]. Such nonintegrable systems are commonly expected to have vanishing Drude weights [30] and potentially exhibit diffusive transport, e.g., due to quantum chaos. Notably, the onset of diffusive hydrodynamics under chaotic quantum dynamics has also been substantiated by exact results obtained in random circuit models [48–50]. However, since the dynamics of real interacting systems with many degrees of freedom poses a formidable challenge to theory and numerics, signatures of clean diffusion have been found only for selected examples [34,51–54].

jonasrichter@uos.de

[†]rsteinig@uos.de

In this context, we study the dynamics of magnetization and energy in a nonintegrable spin-1/2 system with ladder geometry. In particular, we will demonstrate an efficient numerical approach based on the concept of typicality [22–25,55–60], which enables us to study large system sizes and to detect various signatures of diffusive transport.

This paper is structured as follows. We introduce the model in Sec. II and define the observables in Sec. III. In Sec. IV, we outline our numerical approach and present our results in Sec. V. We summarize and conclude in Sec. VI.

II. THE MODEL

The Hamiltonian $\mathcal{H} = \sum_{l=1}^{L} h_l$ of the spin-1/2 ladder with periodic boundary conditions reads

$$h_{l} = J_{||} \sum_{k=1}^{2} \mathbf{S}_{l,k} \cdot \mathbf{S}_{l+1,k} + \frac{J_{\perp}}{2} \sum_{n=l}^{l+1} \mathbf{S}_{n,1} \cdot \mathbf{S}_{n,2}, \qquad (1)$$

where $\mathbf{S}_{\mathbf{r}} = (S_{\mathbf{r}}^x, S_{\mathbf{r}}^y, S_{\mathbf{r}}^z)$ are spin-1/2 operators on lattice site $\mathbf{r} = (l, k)$, $J_{\parallel} = 1$ denotes the coupling along the legs (and sets the energy scale throughout this paper), and $J_{\perp} \ge 0$ is the coupling on the *L* rungs. While, for $J_{\perp} = 0$, \mathcal{H} decouples into two separate chains and is integrable in terms of the Bethe ansatz [61], this integrability is broken for any $J_{\perp} \ne 0$. Numerous works [62–67] have explored the dynamics of the spin ladder (1), including various modifications such as four-spin terms [68], Kitaev-type couplings [69], and XX-ladder systems [70,71]. However, while these studies often discuss either spin *or* energy transport, they also mostly focus exclusively on the dynamics of densities *or* currents, either in time *or* in frequency.

In this paper, we do not focus on a particular quantity and representation and provide a comprehensive picture of high-temperature dynamics in the spin-1/2 ladder. As a main result, we unveil the existence of genuine diffusion for *both* magnetization and energy. In particular, this result is based on *the combination of four distinct signatures*: (i) Gaussian density profiles, (ii) time-independent diffusion coefficients, (iii) exponentially decaying density modes, and (iv) Lorentzian line shapes of the dynamical structure factor. We present these signatures for large systems with up to 40 lattice sites.

III. SETUP AND OBSERVABLES

We study the dynamics of time-dependent expectation values

$$p_{l}(t) = \langle \psi(t) | \varrho_{l} | \psi(t) \rangle, \quad \varrho_{l} = \begin{cases} S_{l,1}^{z} + S_{l,2}^{z}, \\ h_{l}, \end{cases}$$
(2)

where the time argument has to be understood as $|\psi(t)\rangle = e^{-i\mathcal{H}t} |\psi(0)\rangle$ and the operator ϱ_l denotes the local densities of magnetization or energy (see Fig. 1). Furthermore, the (unnormalized) pure initial state $|\psi(0)\rangle$ is prepared as

$$|\psi(0)\rangle = \frac{\sqrt{\tilde{\varrho}_{L/2}} |\varphi\rangle}{\sqrt{\langle \varphi |\varphi\rangle}}; \quad |\varphi\rangle = \sum_{k=1}^{d} c_k |\varphi_k\rangle, \qquad (3)$$

where the complex coefficients c_k are randomly drawn from a Gaussian distribution with zero mean (Haar measure [72]) and $|\varphi_k\rangle$ denote a set of orthonormal basis states (e.g., the



FIG. 1. Sketch of the local densities ϱ_l . (a) Local magnetization $S_{l,1}^z + S_{l,2}^z$. (b) Local energy h_l . Note that h_l is defined with $J_{\perp}/2$ [see Eq. (1)].

Ising basis) of the full Hilbert space with dimension $d = 4^L$. Moreover, the operator $\tilde{\varrho}_{L/2} = \varrho_{L/2} + c$ in Eq. (3) is essentially equivalent to $\varrho_{L/2}$ except for a constant offset which renders the eigenvalues of $\tilde{\varrho}_{L/2}$ nonnegative. Exploiting the concept of dynamical typicality [72,73], as well as $\text{Tr}[\varrho_l] = 0$, the expectation value $p_l(t)$ can be connected to an equilibrium correlation function at formally infinite temperature (see Sec. IV),

$$p_l(t) = \langle \varrho_l(t)\varrho_{L/2} \rangle + \epsilon, \qquad (4)$$

with $\langle \cdot \rangle = \text{Tr}[\cdot]/d$. Note that the statistical error $\epsilon = \epsilon(|\varphi\rangle)$ scales as $\epsilon \propto 1/\sqrt{d}$ and is negligibly small for all system sizes studied here [55,57,72,73].

In addition to the spatiotemporal correlation functions (4), the respective correlations in momentum space can be obtained by a lattice Fourier transform [74],

$$p_q(t) = \sum_l e^{iq(l-L/2)} p_l(t) = \langle \varrho_q(t) \varrho_{-q} \rangle, \tag{5}$$

where translational invariance has been exploited and $\varrho_q = \sqrt{1/L} \sum_l e^{iql} \varrho_l$, with discrete momenta $q = 2\pi k/L$ and $k = 0, 1, \dots, L-1$. Furthermore, a subsequent Fourier transform from the time to frequency domain eventually yields the dynamical structure factor $S(q, \omega)$,

$$p_q(\omega) = \int_{-t_{\text{max}}}^{t_{\text{max}}} e^{i\omega t} p_q(t) \, dt = S(q, \omega), \tag{6}$$

with the finite frequency resolution $\delta \omega = \pi / t_{\text{max}}$.

It is further instructive to establish a relation between density dynamics and current-current correlation functions. To this end, let us introduce the time-dependent diffusion coefficient [75]

$$D(t) = \frac{1}{\chi} \int_0^t \frac{\langle j(t')j\rangle}{L} dt',$$
(7)

where $\chi = \lim_{q \to 0} \langle \varrho_q \varrho_{-q} \rangle$ denotes the isothermal susceptibility [76] and the spin- or energy-current operators $j = \sum_l j_l$ follow from the lattice continuity equation $\partial_t \varrho_l = i[\mathcal{H}, \varrho_l] = j_{l-1} - j_l$. More details on current operators and autocorrelation functions are provided in Appendix A.

To proceed, we note that the states $|\psi(0)\rangle$ realize an initial density profile $p_l(0)$ which exhibits a δ peak at l = L/2 [77]. This initial δ peak will gradually broaden with time, and its spatial variance for $t \ge 0$ is given by

$$\sigma(t)^{2} = \sum_{l=1}^{L} l^{2} \delta p_{l}(t) - \left(\sum_{l=1}^{L} l \delta p_{l}(t)\right)^{2}, \qquad (8)$$

with $\delta p_l(t) = p_l(t) / \sum_l p_l(0)$ and $\sum_l \delta p_l(t) = 1$. Due to the typicality relation in Eq. (4), this variance can be directly connected to the aforementioned diffusion coefficient [78,79],

$$\frac{d}{dt}\sigma(t)^2 = 2D(t).$$
(9)

Note that a diffusive process requires D(t) = D = const, such that $\sigma(t) \propto \sqrt{t}$ [34], above the mean free time.

IV. NUMERICAL APPROACH

A. Dynamical quantum typicality

First, let us derive the typicality relation given in Eq. (4). To this end, we start with an infinite-temperature correlation function,

$$\mathcal{C}_{l,l'}(t) = \frac{\operatorname{Tr}[\varrho_l(t)\tilde{\varrho}_{l'}]}{d} = \frac{\operatorname{Tr}[\varrho_l(t)\varrho_{l'}] + c\operatorname{Tr}[\varrho_l]}{d}, \quad (10)$$

where $\tilde{\varrho}_{l'} = \varrho_{l'} + c$ has a nonnegative spectrum. We realize that Eq. (10) can be simplified if either c = 0 or $\text{Tr}[\varrho_l] = 0$. Focusing on these cases, we find

$$\mathcal{C}_{l,l'}(t) = \frac{\operatorname{Tr}[\varrho_l(t)\varrho_{l'}]}{d} = \langle \varrho_l(t)\varrho_{l'} \rangle, \tag{11}$$

that is, the correlation functions $\langle \varrho_l(t)\tilde{\varrho}_{l'}\rangle$ and $\langle \varrho_l(t)\varrho_{l'}\rangle$ are equivalent. This is the first important observation. Furthermore, exploiting the cyclic invariance of the trace, Eq. (10) can be written as

$$C_{l,l'}(t) = \frac{\operatorname{Tr}[\sqrt{\tilde{\varrho}_{l'} \varrho_l(t)} \sqrt{\tilde{\varrho}_{l'}}]}{d},$$
(12)

where the square-root operation has to be understood in a representation where $\tilde{\varrho}_{l'}$ is diagonal (see Sec. IV B). Using the concept of quantum typicality [22–25,55–60], the trace in Eq. (12) can be replaced by a scalar product with a single pure state $|\varphi\rangle$ which is randomly drawn according to the unitary invariant Haar measure,

$$C_{l,l'}(t) \approx \frac{\langle \varphi | \sqrt{\tilde{\varrho}_{l'}} \varrho_l(t) \sqrt{\tilde{\varrho}_{l'}} | \varphi \rangle}{\langle \varphi | \varphi \rangle} + \epsilon(|\varphi\rangle)$$
(13)

$$= \langle \psi(0) | \varrho_l(t) | \psi(0) \rangle = \langle \psi(t) | \varrho_l | \psi(t) \rangle, \qquad (14)$$

where we have used the definition of $|\psi(0)\rangle$ from Eq. (3) and interpreted the time dependence as a property of the states and not of the operator. The combination of Eqs. (10) to (14) yields the typicality relation (4), where we have chosen l' = L/2without loss of generality.

B. Construction of initial states

Concerning the construction of the pure state $|\psi(0)\rangle$ in Eq. (3), i.e., the evaluation of the square root $\sqrt{\tilde{\varrho}_{L/2}}$, the following comments are in order. On the one hand, in the case of $\varrho_l = S_{l,1}^z + S_{l,2}^z$ this procedure is rather simple since $S_{l,1}^z + S_{l,2}^z$ is naturally diagonal in the Ising basis, which is routinely used as our working basis. On the other hand, in the case of the local energy, $\varrho_l = h_l$ is not diagonal immediately. While this situation usually requires diagonalization, a complete diagonalization of h_l can still be avoided since h_l is a local operator acting nontrivially only on a small part of the product space. Thus, although the preparation of $|\psi(0)\rangle$ becomes more

demanding in the case of $\varrho_l = h_l$, it certainly remains feasible and yields a powerful numerical approach as well. If one still wants to refrain from such square-root constructions, it is also possible to use *two* auxiliary pure states instead of just *one* (see Appendix A). It should be noted, however, that the approach presented in this paper, using just a single pure state, will generally be more favorable concerning memory requirements and run time (even if the initial preparation of $|\psi(0)\rangle$ is costlier).

C. Pure-state propagation

Relying on the typicality relation (4), we calculate spatiotemporal correlation functions for spin and energy densities. The main advantage of this approach comes from the fact that the action of $e^{-i\mathcal{H}t}$ on the pure state $|\psi(0)\rangle$ can be efficiently evaluated *without* the diagonalization of \mathcal{H} , e.g., by means of a Trotter decomposition [80] (see also Appendix B) or by other approaches [81-83]. Moreover, let us stress that the numerical costs of the Fourier transforms (5) and (6) are practically negligible. Therefore, we essentially obtain all information on the dynamics of either magnetization or energy from the time evolution of the *single* pure state $|\psi(t)\rangle$ and the measurement of L local operators ρ_l (see Appendix C). In practice, this pure-state approach enables us to treat ladders with up to L = 20 rungs, i.e., 40 lattice sites in total. If not stated otherwise, we always take into account the full Hilbert space (e.g., $d \approx 10^{12}$ for L = 20).

D. Finite temperatures

Although the focus of this paper is on quantum dynamics at formally infinite temperature, let us briefly outline how finite-temperature correlations can be obtained based on purestate calculations as well. On the one hand, a straightforward approach is the construction of the typical pure state $|\varphi_{\beta}\rangle$ according to [57,60]

$$|\varphi_{\beta}\rangle = e^{-\beta \mathcal{H}/2} |\varphi\rangle, \qquad (15)$$

where the reference pure state $|\varphi\rangle$ corresponds to infinite temperature and has been introduced in Eq. (3). Analogous to the real-time evolution, the action of $e^{-\beta \mathcal{H}/2}$ can be efficiently evaluated by an iterative forward propagation, but now in imaginary time. More details on finite-temperature calculations using $|\varphi_{\beta}\rangle$ can be found in Appendix A.

On the other hand, another useful class of pure states $|\tilde{\varphi}_{\alpha,\beta}\rangle$, which was put forward in Ref. [84], is constructed as

$$|\tilde{\varphi}_{\alpha,\beta}\rangle \propto e^{-\beta(\mathcal{H}-\alpha\varrho_{l'})/2} |\varphi\rangle.$$
 (16)

In particular, depending on the size of the control parameter α , these states can be used not only to obtain linear-response correlation functions but also to calculate far-from-equilibrium quantum dynamics [21,84].

Finally, let us note that, since the effective Hilbert-space dimension shrinks for $\beta > 0$, the statistical error ϵ of the typicality approximation [see Eq. (4)] is generally larger than the infinite-temperature limit. Nevertheless, even for $\beta > 0$ this error still decreases exponentially with system size, and accurate calculations remain possible for moderate temperatures.



FIG. 2. Real-time broadening of density profiles $p_l(t)$ for spin $\varrho_l = S_{l,1}^z + S_{l,2}^z$ and energy $\varrho_l = h_l$. The other parameters are $J_{\parallel} = J_{\perp} = 1$ and L = 20.

V. RESULTS

We now present our numerical results. In Sec. V A, we start with the analysis of density dynamics in real space. The corresponding structure factors are then discussed for magnetization in Sec. V B and for energy in Sec. V C.

A. Real-space dynamics

To begin with, let us focus on the isotropic case $J_{\parallel} = J_{\perp} = 1$. Starting with dynamics in time and real space, Figs. 2(a) and 2(b) show the density profiles $p_l(t)$ of magnetization and energy for large systems with L = 20. One can clearly observe the initial δ peak at t = 0 (or almost δ peak [77]), which broadens for times t > 0. Moreover, on the (short) timescales depicted, $p_l(t)$ does not reach the boundaries of the system; that is, trivial finite-size effects do not occur.

For a more detailed discussion, Figs. 3(a) and 3(b) show cuts of $p_l(t)$ at fixed times t = 0, 1, 2, 4. For these times, one finds that the data are well described by Gaussians over



several orders of magnitude,

$$p_l(t) \propto \exp\left[-\frac{(l-L/2)^2}{2\sigma(t)^2}\right].$$
 (17)

While these Gaussians already suggest diffusion for both magnetization and energy (see Appendix D), it is only a sufficient criterion if $\sigma(t)$ scales as $\sigma(t) \propto \sqrt{t}$ as well. Consequently, Figs. 3(c) and 3(d) show the widths of the density profiles obtained from Eq. (8) (symbols) in comparison with the respective quantities D(t) and $\sigma(t)$ (lines), calculated from the current autocorrelations [see Eqs. (7) and (9)]. Generally, one observes excellent agreement between density and current dynamics. Moreover, after a linear increase at short times, D(t) eventually saturates at a constant plateau $D(t) \approx \text{const}$, and correspondingly, $\sigma(t) \propto \sqrt{t}$. Thus, based on our numerical analysis in time and real space, we unveil the existence of diffusive transport in the spin-1/2 ladder for both magnetization and energy. This is the first main result of this paper.

Let us now briefly present magnetization profiles also for other ratios $J_{\perp}/J_{\parallel} \neq 1$. To this end, Fig. 4 shows the real-space density profiles $p_l(t)$ at fixed times t = 1, 2, 4 for interchain couplings $J_{\perp} = 0.5, 1, 2$ ($J_{\parallel} = 1$). Generally, we find that the profiles are very similar to each other for all t and J_{\perp} shown here. In particular, all profiles are convincingly described by Gaussian fits over several orders of magnitude.

B. Spin structure factor

Next, let us also study magnetization dynamics in momentum space, where the lattice diffusion equation decouples into separate Fourier modes (see Appendix D).

1. Long-wavelength limit

In Fig. 5(a), the density modes $p_q(t)$ for $J_{\parallel} = J_{\perp} = 1$ are shown for various momenta q in a semilogarithmic plot. On the one hand, for large $q = \pi$, $p_q(t)$ exhibits pronounced oscillations and essentially decays on a timescale $t \sim 5$. On the other hand, for the two smallest wave numbers k = 1 and k = 2, we find clean exponential relaxation

$$p_q(t) \propto e^{-\tilde{q}^2 D t},$$
 (18)

FIG. 3. (a) and (b) Density profiles $p_l(t)$ for spin and energy at fixed times t = 0 (δ peak) and t = 1, 2, 4 (arrow). The dashed lines are Gaussian fits to the data. (c) and (d) Widths of the density profiles (symbols) obtained from Eq. (8), as well as D(t) and $\sigma(t)$ (lines) calculated from current autocorrelations [see Eqs. (7) and (9)]. For sufficiently long times, one finds $D(t) \approx$ const. The other parameters are $J_{\parallel} = J_{\perp} = 1, L = 20$ (densities), L = 18 (spin current), and L = 15 (energy current [67]).



FIG. 4. Magnetization profiles $p_l(t)$ at fixed times t = 1, 2, 4. Comparison between different interchain couplings $J_{\perp} = 0.5, 1, 2$. The dashed lines are Gaussian fits to the data. The other parameters are $J_{\parallel} = 1$ and L = 18.

where we have introduced the abbreviation $\tilde{q}^2 = 2[1 - \cos(q)] \approx q^2$ for sufficiently small q and D = const can be extracted from the constant plateau in Fig. 3(c). Going from the time to frequency domain, Fig. 5(b) shows the corresponding dynamical structure factors $p_q(\omega)$ for k = 1, 2. One observes



FIG. 5. Spin dynamics. (a) $p_q(t)$ for wave numbers k = 1, 2, 8in a semilogarithmic plot. Lines are exponential functions according to Eq. (18). (b) $p_q(\omega)$ for k = 1, 2. Lines are Lorentzians according to Eq. (19). (c) $p_q(\omega)$ for k = 8. Comparison with the XY model [Eq. (20)] and an effective model (γ XY) where the original memory kernel of the Bessel function is exponentially damped (here $\gamma = 0.6$) [85]. The XY data are multiplied by a factor of 2 in order to account for the two legs of the ladder. (d) $p_q(\omega)$ in the full Brillouin zone. Arrows show the data of (b) and (c). Other parameters are $J_{\parallel} = J_{\perp} =$ 1, L = 16, and $\delta\omega = \pi/150$.



FIG. 6. Spin dynamics for different interchain couplings $J_{\perp} = 0.5, 1, 2.$ (a) $p_q(t)$ for the two smallest wave numbers, k = 1, 2. (b) and (c) $p_q(\omega)$ for k = 1 and k = 9, respectively. Other parameters are $J_{\parallel} = 1, L = 18$, and $\delta \omega = \pi/25$.

that the data can be accurately described by Lorentzians of the form

$$p_q(\omega) \propto \frac{1}{\omega^2 + \tilde{q}^4 D^2}.$$
 (19)

Note that, while we display the time data in Fig. 5(a) only up to intermediate timescales, the Fourier transform (6) is routinely performed for a much longer cutoff time (here $t_{max} = 150$) in order to achieve a high frequency resolution. The exponential relaxation [Fig. 5(a)] and the Lorentzian line shapes [Fig. 5(b)] clearly confirm our earlier observations in the context of Fig. 3, i.e., the occurrence of genuine spin diffusion in the spin-1/2 ladder. This is another main result of the present work.

Eventually, Fig. 5(d) shows the dynamical structure factor $p_q(\omega)$ for all momenta q. Let us stress again that within our numerical approach the calculation of these density modes essentially does not require any additional resources. On the one hand, for small momenta $q \rightarrow 0$, one can clearly identify the diffusion peaks discussed above. On the other hand, in the center of the Brillouin zone, we find that $p_q(\omega)$ exhibits a broad continuum. This short-wavelength limit will be discussed below in more detail.

In order to corroborate once more that the emergence of diffusive transport is not restricted to the isotropic point, Fig. 6 shows the structure factors $p_q(t)$ and $p_q(\omega)$ for $J_{\perp} = 0.5$, 1, 2. For the smallest wave number k = 1, we find a clean exponential decay of $p_q(t)$ with a decay rate $-\tilde{q}^2 D$, which is almost identical for all J_{\perp} shown here. This fact is also reflected in the Lorentzian shape of $p_q(t)$ for this momentum [Fig. 6(b)], which essentially coincides for all strengths of interchain couplings. On the other hand, for wave number k = 2, we find that $p_q(t)$ for $J_{\perp} = 0.5$ shows some deviations from an exponential; that is, the hydrodynamic regime becomes smaller for smaller J_{\perp} , which can be explained by the increased mean free path of spin excitations.

2. Short-wavelength limit

In addition to the long-wavelength limit, Fig. 5(c) shows $p_q(\omega)$ at momentum $q = \pi$. For this momentum, one finds that $p_q(\omega)$ is practically ω independent up to $\omega \leq 2$ and



FIG. 7. Comparison of $p_{q=\pi}(t)$ between the spin ladder with $J_{\parallel} = J_{\perp} = 1$ [Fig. 5(a)], the bare XY model (Bessel function), and the effective dynamics $\tilde{p}_q(t)$ (γ XY) [see Eqs. (21) and (23)] with damping $\gamma = 0.6$. The XY data are multiplied by a factor of 2 in order to account for the two legs of the ladder.

exhibits a constant plateau. It is instructive to compare this result to the dynamics of the one-dimensional XY model. Since the XY chain is equivalent to a model of free fermions, $p_q(\omega)$ is known exactly [74] and reads (for $\beta \to 0$, $J_{\parallel} = 1$, and $q = \pi$)

$$p_{q=\pi}(\omega) = \int_{-\infty}^{\infty} e^{i\omega t} \frac{\mathcal{J}_0(2t)}{4} dt = \frac{\Theta(2-|\omega|)}{2\sqrt{4-\omega^2}}.$$
 (20)

Here, $\mathcal{J}_0(t)$ is the Bessel function of the first kind (and order zero), and $p_{q=\pi}(\omega)$ exhibits a square-root divergence at $\omega = 2$ [see Fig. 5(c)]. Following an approach introduced in Ref. [85], the dynamics $p_q(t) \propto \mathcal{J}_0(2t)$ is interpreted as being generated by an integro-differential equation comprising a *memory kernel K*(*t*),

$$\frac{d}{dt} p_q(t) = -\int_0^t K(t - t') p_q(t') dt'.$$
 (21)

Equation (21) establishes a direct correspondence between $p_q(t)$ and $K(\tau)$ and can be evaluated in both directions. Thus, given the original dynamics $p_q(t) \propto \mathcal{J}_0(2t)$, the respective memory kernel $K(\tau)$ can be calculated, e.g., numerically. In fact, given the expression in Eq. (20), $K(\tau)$ can even be obtained analytically and reads

$$K(\tau) = \frac{2\mathcal{J}_1(2\tau)}{\tau}.$$
 (22)

Comparing the bare XY model with the full spin ladder (1), the additional rung couplings and the $S_{\mathbf{r}}^z S_{\mathbf{r}'}^z$ terms are treated as a perturbation giving rise to an exponential damping of this memory kernel (for small perturbations) [85,86],

$$\tilde{K}(\tau) = K(\tau)e^{-\gamma\tau} .$$
(23)

This new memory kernel $\tilde{K}(\tau)$ is then used to numerically evaluate Eq. (21) in order to obtain the new (damped) dynamics $\tilde{p}_q(t)$. As shown in Fig. 5(c), the effective dynamics generated by this (heuristic) approach with $\gamma = 0.6$ reproduces the structure factor of the spin ladder remarkably well, even though the perturbation is not small. Moreover, as shown in Fig. 7, this convincing agreement between spin ladder and effective model can be observed not only in frequency space but also in real time. Let us note that, while $\gamma = 0.6$ is found to describe the data most accurately, a more quantitative



FIG. 8. Energy dynamics. (a) $p_q(t)$ for k = 1, 2, 8. (b) $p_q(\omega)$ for k = 1, 2. (c) $p_q(\omega)$ for k = 7, 8. (d) $p_q(\omega)$ in the full Brillouin zone. Other parameters are $J_{\parallel} = J_{\perp} = 1, L = 16$, and $\delta \omega = \pi/150$.

understanding of this specific value goes beyond the scope of the present paper (see also Ref. [86]).

Thus, while clear signatures of diffusion can be found for long-wavelength modes [see Figs. 5(a) and 5(b)], the relaxation of $p_q(t)$ for $q \rightarrow \pi$ can be qualitatively understood as the (damped) dynamics of free fermions, e.g., since the wavelength is smaller than the mean free path. Note that similar behavior has also been found for XXZ chains [74,87].

C. Energy structure factor

We now present results for energy dynamics in momentum space. Analogously to the discussion in the context of Fig. 5, Fig. 8 shows the energy structure factors $p_q(t)$ and $p_q(\omega)$ of the isotropic spin ladder for short and long wavelengths. Generally, the results for energy dynamics are very similar to the previously discussed case of magnetization; that is, $p_q(t)$ decays exponentially for small q, while the corresponding $p_q(\omega)$ exhibits a Lorentzian line shape. Thus, the data shown in Fig. 8 confirm the existence of diffusive energy transport as well (see also Refs. [62,66,67]).

Finally, it is instructive to discuss the effect of an additional uniform magnetic field B > 0 in the z direction; that is, the new local energy \tilde{h}_l reads

$$\tilde{h}_{l} = h_{l} + \frac{B}{2} \sum_{n=l}^{l+1} \sum_{k=1}^{2} S_{n,k}^{z}, \qquad (24)$$

where h_l is defined according to Eq. (1). Such a modification results in a magnetothermal correction, and the heat current takes on the form

$$\tilde{j}_E = j_E + B j_S, \tag{25}$$

where the spin current j_S is independent of *B*. For specific expressions of j_S and j_E , see Appendix A. In Fig. 9, we depict the energy structure factors $p_q(t)$ and $p_q(\omega)$ in the time as well as the frequency domain for various momenta *q*.



FIG. 9. Energy dynamics in the presence of a magnetic field. Calculations are performed only in the subsector with $S^z = 0$. The other parameters are $J_{\parallel} = J_{\perp} = B = 1$, L = 14, and $\delta \omega = \pi/25$.

Note that we restrict ourselves to the symmetry subspace with magnetization $S^z = 0$. Generally, we find that the presence of a finite magnetic field does not qualitatively change the behavior of $p_q(t)$ and $p_q(\omega)$. Again, one can observe an exponential decay for the two smallest wave numbers k = 1, 2 and, correspondingly, a Lorentzian line shape in frequency space at these momenta. Compared to the results with B = 0 shown in Fig. 8, one might even argue that those signatures of diffusion are slightly improved due to the finite magnetic field.

VI. CONCLUSION

To summarize, we have studied spin and energy dynamics in the spin-1/2 ladder for large systems with up to 40 lattice sites. Our state-of-the-art numerical simulations have unveiled the existence of genuine diffusion for both spin and energy. In particular, this finding is based on four distinct signatures which have all been equally well detected: (i) Gaussian density profiles, (ii) time-independent diffusion coefficients, (iii) exponentially decaying density modes, and (iv) Lorentzian line shapes of the dynamical structure factor. Combining (i)-(iv), this paper provides a comprehensive picture of hightemperature dynamics in the spin-1/2 ladder. Promising directions of research include, e.g., the application of the pure-state approach to a larger class of condensed-matter systems and a wider range of temperatures [88]. In this context, whether the observed signatures of diffusion persist at lower temperatures is also an intriguing question.

ACKNOWLEDGMENTS

This work has been funded by the Deutsche Forschungsgemeinschaft (DFG) - Grants No. 397107022 (GE 1657/3-1), No. 397067869 (STE 2243/3-1), No. 355031190 - within the DFG Research Unit FOR 2692. Additionally, we gratefully acknowledge the computing time granted by the JARA-HPC Vergabegremium and provided on the JARA-HPC Partition of the supercomputer JUWELS at Forschungszentrum Jülich. J.H. was supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Materials Sciences and Engineering Division.

APPENDIX A: CURRENT OPERATORS AND AUTOCORRELATIONS

The spin current j_s is defined as [66]

$$j_{S} = J_{\parallel} \sum_{l=1}^{L} \sum_{k=1}^{2} \left(S_{l,k}^{x} S_{l+1,k}^{y} - S_{l,k}^{y} S_{l+1,k}^{x} \right).$$
(A1)

Moreover, the energy current $j_E = j_{\parallel} + j_{\perp}$ can be written as the sum of a longitudinal part and a perpendicular part, which read [62,67]

$$j_{\parallel} = J_{\parallel}^2 \sum_{l=1}^{L} \sum_{k=1}^{2} \mathbf{S}_{l-1,k} \cdot (\mathbf{S}_{l,k} \times \mathbf{S}_{l+1,k}), \qquad (A2)$$

$$j_{\perp} = \frac{J_{\parallel}J_{\perp}}{2} \sum_{l=1}^{L} \sum_{k=1}^{2} (\mathbf{S}_{l-1,k} - \mathbf{S}_{l+1,k}) \cdot (\mathbf{S}_{l,k} \times \mathbf{S}_{l,3-k}).$$
(A3)

In order to calculate current-current correlation functions $\langle j(t)j \rangle$ by means of a typicality-based approach directly (at finite or infinite temperature), we use the two auxiliary pure states [60],

$$|\varphi(\beta,t)\rangle = e^{-i\mathcal{H}t}e^{-\beta\mathcal{H}/2} |\varphi\rangle, \qquad (A4)$$

$$|\phi(\beta,t)\rangle = e^{-i\mathcal{H}t} j e^{-\beta\mathcal{H}/2} |\varphi\rangle, \qquad (A5)$$

which differ by only the additional current operator in Eq. (A5), where $|\varphi\rangle$ is again a random state drawn according to the Haar measure [see Eq. (3)]. Then, we can write [60]

$$\langle j(t)j\rangle = \frac{\langle \varphi(\beta,t)| \, j \, |\phi(\beta,t)\rangle}{\langle \varphi(\beta,0)|\varphi(\beta,0)\rangle} + \epsilon(|\varphi\rangle), \tag{A6}$$

where again $\epsilon(|\varphi\rangle) \propto 1/\sqrt{d}$ for $\beta \to 0$. Of course, by replacing the current operator *j* in Eqs. (A5) and (A6), it is straightforward to generalize the above approach in order to calculate dynamic correlation functions also for other operators.

While we already introduced the time-dependent diffusion coefficient D(t) for spin and energy transport in Eq. (7), the respective ac conductivities at finite frequency ω are given by the Fourier transform of the current autocorrelations,

$$\operatorname{Re}\sigma(\omega) = \frac{1 - e^{-\beta\omega}}{\omega L} \operatorname{Re} \int_0^\infty e^{i\omega t} \langle j_S(t) j_S \rangle \, dt, \qquad (A7)$$

$$\operatorname{Re}\kappa(\omega) = \beta \frac{1 - e^{-\beta\omega}}{\omega L} \operatorname{Re} \int_0^\infty e^{i\omega t} \langle j_E(t) j_E \rangle \, dt. \quad (A8)$$

Note that the spin conductivity $\sigma(\omega)$ from Eq. (A7) must not be confused with the spatial variance $\sigma(t)$ introduced in Eq. (8). Omitting the possibility of finite Drude weights, the corresponding dc conductivities are given by $\lim_{\omega\to 0} \sigma[\kappa](\omega) = \sigma[\kappa]_{dc}$ and can be connected to the diffusion constant via the Einstein relation $D = D(t \to \infty) =$ $\sigma[\kappa]_{dc}/\chi$ [see Eq. (7)]. Let us note that, since the Fourier transforms in Eqs. (A7) and (A8) can, in practice, be evaluated only up to a finite cutoff time $t_{max} < \infty$, the frequency



FIG. 10. Current autocorrelations. (a) Spin current j_S for $J_{\perp} = 0.5$, 1, 2 and $\beta = 0$. (b) Energy current j_E for $J_{\perp} = 1$, 2 and $\beta = 0$. (c) Spin and energy current for $\beta = 1$ and $J_{\perp} = 1$. For comparison, we depict tDMRG data digitized from Ref. [66]. Note that in the case of j_E we restrict ourselves to the symmetry subspaces with momentum k = 0. Other parameters are $J_{\parallel} = 1$, L = 13 (spin), and L = 15 (energy).

resolution of $\sigma[\kappa](\omega)$ is finite as well [see also Eq. (6) in the main text].

In Figs. 10(a) and 10(b), the current autocorrelations $\langle j(t)j \rangle$ at $\beta = 0$ are shown for spin and energy transport, respectively. While in Fig. 10(a) we show data for smaller systems with only L = 13 rungs, the energy current in Fig. 10(b) is calculated for systems with L = 15. Note, however, that in the latter case, we restrict ourselves to the symmetry subspaces with momentum k = 0 as the current is known to be essentially independent of k for such system sizes. [This crystal momentum k should not be confused with the wave number k below Eq. (5).] In all cases shown here, we observe that $\langle i(t)i \rangle$ decays to approximately zero, consistent with the absence of ballistic transport in a nonintegrable system. In the case $J_{\perp}/J_{\parallel} = 1$, our results are additionally compared to data digitized from Ref. [66] obtained by a time-dependent density matrix renormalization group (tDMRG) approach. Generally, one finds convincing agreement between both methods; that is, our data for L = 13, 15 are free of significant finite-size effects and represent the thermodynamic limit. In Fig. 10(c), spin and energy autocorrelations are depicted for the finite temperature $\beta = 1$. Also in this case, we observe that the purestate method accurately reproduces the tDMRG data. Thus,



FIG. 11. (a) Spin conductivity $\sigma(\omega)$ for $J_{\perp} = 0.5, 1, 2$. (b) Energy conductivity $\kappa(\omega)$ for $J_{\perp} = 1, 2$. We show data for two different frequency resolutions, $\delta\omega = \pi/10$ (solid symbols) and $\delta\omega = \pi/50$ (open symbols). For comparison, tDMRG data digitized from Ref. [66] are shown. In (b), we also compare our results to data obtained from the microcanonical Lanczos method (MCLM) [62]. In the case of $\kappa(\omega)$ we restrict ourselves to the symmetry subspaces with momentum k = 0. Other parameters are $J_{\parallel} = 1, \beta = 0, L = 13$ (spin), and L = 15 (energy).

typical pure states yield an efficient approach to correlation functions at finite temperatures as well.

In Figs. 11(a) and 11(b) the respective ac conductivities $\sigma(\omega)$ and $\kappa(\omega)$ at $\beta = 0$ are shown, i.e., the Fourier transforms of the data in Figs. 10(a) and 10(b). Particularly, we compare data with two different frequency resolutions, $\delta \omega = \pi/10$ and $\delta \omega = \pi/50$, i.e., a rather short and a significantly longer cutoff time t_{max} in Eqs. (A7) and (A8). In all cases, we observe a well-behaved dc conductivity $\sigma[\kappa]_{\text{dc}} > 0$ which does not (significantly) depend on t_{max} , except $\kappa(\omega)$ for $J_{\perp} = 2$. Moreover, our data are again in good agreement with existing data obtained by tDMRG [66] and by the microcanonical Lanczos method [62].

APPENDIX B: TROTTER DECOMPOSITION

Let us briefly give some details on the time evolution of pure quantum states by means of a Trotter decomposition. To begin with, we note that the time-dependent Schrödinger equation

$$i\partial_t |\psi(t)\rangle = \mathcal{H} |\psi(t)\rangle \tag{B1}$$

is formally solved by

$$|\psi(t')\rangle = U(t,t') |\psi(t)\rangle, \qquad (B2)$$

with $U(t, t') = e^{-i\mathcal{H}(t'-t)}$, where we have set $\hbar = 1$. While the exact evaluation of Eq. (B2) requires the diagonalization of \mathcal{H} , we here approximate the time-evolution operator U(t, t') by means of a Trotter product formula. Specifically, a second-

order approximation of $U(t, t + \delta t) = U(\delta t)$ is given by

$$\widetilde{U}_{2}(\delta t) = e^{-i\frac{\delta t}{2}\mathcal{H}_{k}} \cdots e^{-i\frac{\delta t}{2}\mathcal{H}_{1}} e^{-i\frac{\delta t}{2}\mathcal{H}_{1}} \cdots e^{-i\frac{\delta t}{2}\mathcal{H}_{k}}, \qquad (B3)$$

where $\mathcal{H} = \mathcal{H}_1 + \cdots + \mathcal{H}_k$. This approximation is then bounded by

$$||U(\delta t) - \widetilde{U}_2(\delta t)|| \ll c_2 \,\delta t^3,\tag{B4}$$

where c_2 is a positive constant. In practice, the Hamiltonian is decomposed into the *x*, *y*, and *z* components of the spin operators, i.e., $\mathcal{H} = \mathcal{H}_x + \mathcal{H}_y + \mathcal{H}_z$. Since the computational basis states are eigenstates of the S^z operators, the representation $e^{-i\delta t \mathcal{H}_z}$ is diagonal by construction and only changes the input state by altering the phase of each of the basis vectors. Using an efficient basis rotation in the eigenstates of the S^x or S^y operators, the operators $e^{-i\delta t \mathcal{H}_x}$ and $e^{-i\delta t \mathcal{H}_y}$ can act as $e^{-i\delta t \mathcal{H}_z}$ as well [80].

APPENDIX C: FOURIER TRANSFORMS

Let us comment on the derivation of Eqs. (5) and (6) from the main part of this paper. Referring to Eq. (4), we realize that a cut through the density profile $p_l(t)$ at fixed lattice site *l* is equivalent to the correlation function $\langle \varrho_l(t) \varrho_{L/2} \rangle$. It is now instructive to perform the following calculation:

$$\langle \varrho_q(t)\varrho_{-q}\rangle = \frac{1}{L}\sum_{l=1}^L \sum_{l'=1}^L e^{iq(l-l')} \langle \varrho_l(t)\varrho_{l'}\rangle \tag{C1}$$

$$=\sum_{l=1}^{L}e^{iql}\langle \varrho_{L/2+l}(t)\varrho_{L/2}\rangle \tag{C2}$$

$$=\sum_{l=1}^{L} e^{iq(l-L/2)} p_l(t) = p_q(t),$$
(C3)

where we have exploited translational invariance in order to compress the original double sum. Thus, we find that the intermediate structure factor $\langle \varrho_q(t) \varrho_{-q} \rangle$ can be easily obtained by a lattice Fourier transform of the real-space correlations. Furthermore, this momentum-space correlation function can also be transferred to what is usually referred to as the *dy*-

- D. C. Johnston, R. K. Kremer, M. Troyer, X. Wang, A. Klümper, S. L. Bud'ko, A. F. Panchula, and P. C. Canfield, Phys. Rev. B 61, 9558 (2000).
- [2] A. V. Sologubenko, K. Giannó, H. R. Ott, U. Ammerahl, and A. Revcolevschi, Phys. Rev. Lett. 84, 2714 (2000).
- [3] C. Hess, C. Baumann, U. Ammerahl, B. Büchner, F. Heidrich-Meisner, W. Brenig, and A. Revcolevschi, Phys. Rev. B 64, 184305 (2001).
- [4] B. Lake, D. A. Tennant, J.-S. Caux, T. Barthel, U. Schollwöck, S. E. Nagler, and C. D. Frost, Phys. Rev. Lett. **111**, 137205 (2013).

namical structure factor $S(q, \omega)$ by another Fourier transform from the time to frequency domain,

$$S(q,\omega) = \int_{-\infty}^{\infty} e^{i\omega t} \langle \varrho_q(t) \varrho_{-q} \rangle \, dt \tag{C4}$$

$$\approx \int_{-t_{\text{max}}}^{t_{\text{max}}} e^{i\omega t} p_q(t) \, dt = p_q(\omega), \tag{C5}$$

where the finite cutoff time t_{max} yields a frequency resolution $\delta \omega = \pi / t_{\text{max}}$. Thus, starting from the correlations $p_l(t)$, it is straightforward to also obtain correlation functions in the momentum and frequency domains, which makes our purestate method a rather powerful numerical approach.

APPENDIX D: DIFFUSION IN LATTICE MODELS

In this Appendix, let us discuss in more detail how to detect diffusion in lattice models. To begin with, a process is called diffusive if it fulfills the lattice diffusion equation

$$\frac{d}{dt} p_l(t) = D[p_{l-1}(t) - 2p_l(t) + p_{l+1}(t)],$$
(D1)

with the *time-independent* diffusion constant *D*. In the case of an initial δ peak profile at l = L/2, a specific solution for the time dependence of $p_t(t)$ is given by [54]

$$p_l(t) - p_{eq} = \frac{1}{2} \exp(-2Dt) \mathcal{B}_{l-L/2}(2Dt),$$
 (D2)

where $\mathcal{B}_l(t)$ is the modified Bessel function of the first kind and $p_{eq} = p_{l \neq L/2}(0)$ denotes the homogeneous background. (Note that in this paper we have $p_{eq} = 0$ since $\text{Tr}[\varrho_l] = 0$.) In the case of sufficiently large *L* and long *t*, i.e., if the discrete lattice momenta *q* become dense, this lattice solution can be well approximated by a Gaussian. Such Gaussians are observed in Figs. 3(a) and 3(b). Specifically, the spatial variance of these Gaussians is then also given by $\sigma^2(t) = 2Dt$, i.e., $\sigma(t) \propto \sqrt{t}$.

Given some general density distribution, it is in some cases instructive to study the dynamics in momentum space as well. In this context, a Fourier transform of Eq. (D1) yields

$$\frac{d}{dt} p_q(t) = -2D(1 - \cos q)p_q(t).$$
(D3)

Apparently, the L different Fourier modes $p_q(t)$ are completely decoupled, and their exponentially decaying solutions have already been given in Eq. (18).

- [5] M. Mourigal, M. Enderle, A. Klöpperpieper, J.-S. Caux, A. Stunault, and H. M. Rønnow, Nat. Phys. 9, 435 (2013).
- [6] K. R. Thurber, A. W. Hunt, T. Imai, and F. C. Chou, Phys. Rev. Lett. 87, 247202 (2001).
- [7] H. Maeter, A. A. Zvyagin, H. Luetkens, G. Pascua, Z. Shermadini, R. Saint-Martin, A. Revcolevschi, C. Hess, and H.-H. Klauss, J. Phys.: Condens. Matter 25, 365601 (2013).
- [8] S. A. Wolf, D. D. Awschalom, R. A. Buhrman, J. M. Daughton, S. von Molnár, M. L. Roukes, A. Y. Chtchelkanova, and D. Treger, Science 294, 1488 (2001).
- [9] M. A. Cazalilla and M. Rigol, New J. Phys. 12, 055006 (2010).

- [10] Y. Ikhlef, J. L. Jacobsen, and H. Saleur, Phys. Rev. Lett. 108, 081601 (2012).
- [11] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Rev. Mod. Phys. 83, 863 (2011).
- [12] J. Eisert, M. Friesdorf, and C. Gogolin, Nat. Phys. 11, 124 (2015).
- [13] C. Gogolin and J. Eisert, Rep. Prog. Phys. 79, 056001 (2016).
- [14] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, Adv. Phys. 65, 239 (2016).
- [15] T. Langen, R. Geiger, and J. Schmiedmayer, Annu. Rev. Condens. Matter Phys. 6, 201 (2015).
- [16] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Scholwöck, J. Eisert, and I. Bloch, Nat. Phys. 8, 325 (2012).
- [17] R. Blatt and C. F. Roos, Nat. Phys. 8, 277 (2012).
- [18] J. M. Deutsch, Phys. Rev. A 43, 2046 (1991).
- [19] M. Srednicki, Phys. Rev. E 50, 888 (1994).
- [20] M. Rigol, V. Dunjko, and M. Olshanii, Nature (London) 452, 854 (2008).
- [21] J. Richter, J. Gemmer, and R. Steinigeweg, arXiv:1805.11625.
- [22] J. Gemmer, M. Michel, and G. Mahler, *Quantum Thermody*namics (Springer, Berlin, 2004).
- [23] S. Popescu, A. J. Short, and A. Winter, Nat. Phys. 2, 754 (2006).
- [24] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì, Phys. Rev. Lett. 96, 050403 (2006).
- [25] P. Reimann, Phys. Rev. Lett. 99, 160404 (2007).
- [26] U. Schollwöck, Rev. Mod. Phys. 77, 259 (2005); Ann. Phys. (NY) 326, 96 (2011).
- [27] X. Zotos, F. Naef, and P. Prelovšek, Phys. Rev. B 55, 11029 (1997).
- [28] T. Prosen and E. Ilievski, Phys. Rev. Lett. 111, 057203 (2013).
- [29] L. Vidmar and M. Rigol, J. Stat. Mech. (2016) 064007.
- [30] F. Heidrich-Meisner, A. Honecker, D. C. Cabra, and W. Brenig, Phys. Rev. B 68, 134436 (2003); F. Heidrich-Meisner, A. Honecker, and W. Brenig, Eur. Phys. J. Spec. Top. 151, 135 (2007).
- [31] J. Sirker, R. G. Pereira, and I. Affleck, Phys. Rev. B 83, 035115 (2011).
- [32] M. Žnidarič, Phys. Rev. Lett. 106, 220601 (2011).
- [33] R. Steinigeweg and W. Brenig, Phys. Rev. Lett. 107, 250602 (2011).
- [34] C. Karrasch, J. E. Moore, and F. Heidrich-Meisner, Phys. Rev. B 89, 075139 (2014).
- [35] R. Steinigeweg, F. Jin, D. Schmidtke, H. De Raedt, K. Michielsen, and J. Gemmer, Phys. Rev. B 95, 035155 (2017).
- [36] M. Ljubotina, M. Žnidarič, and T. Prosen, Nat. Commun. 8, 16117 (2017).
- [37] T. Prosen and M. Žnidarič, Phys. Rev. B 86, 125118 (2012).
- [38] C. Karrasch, T. Prosen, and F. Heidrich-Meisner, Phys. Rev. B 95, 060406(R) (2017).
- [39] R. Steinigeweg, F. Jin, H. De Raedt, K. Michielsen, and J. Gemmer, Phys. Rev. E 96, 020105(R) (2017).
- [40] O. A. Castro-Alvaredo, B. Doyon, and T. Yoshimura, Phys. Rev. X 6, 041065 (2016).
- [41] B. Bertini, M. Collura, J. De Nardis, and M. Fagotti, Phys. Rev. Lett. 117, 207201 (2016).
- [42] J. De Nardis, D. Bernard, and B. Doyon, Phys. Rev. Lett. 121, 160603 (2018).
- [43] A. L. Chernyshev and A. V. Rozhkov, Phys. Rev. Lett. 116, 017204 (2016).

- [44] K. R. A. Hazzard, M. van den Worm, M. Foss-Feig, S. R. Manmana, E. G. D. Torre, T. Pfau, M. Kastner, and A. M. Rey, Phys. Rev. A **90**, 063622 (2014).
- [45] C. Karrasch, R. Ilan, and J. E. Moore, Phys. Rev. B 88, 195129 (2013).
- [46] A. Metavitsiadis, X. Zotos, O. S. Barišić, and P. Prelovšek, Phys. Rev. B 81, 205101 (2010).
- [47] J. Herbrych, J. Kokalj, and P. Prelovšek, Phys. Rev. Lett. 111, 147203 (2013).
- [48] C. W. von Keyserlingk, T. Rakovszky, F. Pollmann, and S. L. Sondhi, Phys. Rev. X 8, 021013 (2018).
- [49] A. Nahum, S. Vijay, and J. Haah, Phys. Rev. X 8, 021014 (2018).
- [50] V. Khemani, A. Vishwanath, and D. A. Huse, Phys. Rev. X 8, 031057 (2018).
- [51] M. Michel, G. Mahler, and J. Gemmer, Phys. Rev. Lett. 95, 180602 (2005).
- [52] C. Mejía-Monasterio, T. Prosen, and G. Casati, Europhys. Lett. 72, 520 (2005).
- [53] M. Medenjak, K. Klobas, and T. Prosen, Phys. Rev. Lett. 119, 110603 (2017).
- [54] J. Richter, F. Jin, H. De Raedt, K. Michielsen, J. Gemmer, and R. Steinigeweg, Phys. Rev. B 97, 174430 (2018).
- [55] A. Hams and H. De Raedt, Phys. Rev. E 62, 4365 (2000).
- [56] T. Iitaka and T. Ebisuzaki, Phys. Rev. Lett. 90, 047203 (2003).
- [57] S. Sugiura and A. Shimizu, Phys. Rev. Lett. 111, 010401 (2013).
- [58] T. A. Elsayed and B. V. Fine, Phys. Rev. Lett. 110, 070404 (2013).
- [59] T. Monnai and A. Sugita, J. Phys. Soc. Jpn. 83, 094001 (2014).
- [60] R. Steinigeweg, J. Gemmer, and W. Brenig, Phys. Rev. Lett. 112, 120601 (2014).
- [61] A. Klümper and K. Sakai, J. Phys. A 35, 2173 (2002).
- [62] X. Zotos, Phys. Rev. Lett. 92, 067202 (2004).
- [63] P. Jung, R. W. Helmes, and A. Rosch, Phys. Rev. Lett. 96, 067202 (2006).
- [64] S. Langer, F. Heidrich-Meisner, J. Gemmer, I. P. McCulloch, and U. Schollwöck, Phys. Rev. B 79, 214409 (2009).
- [65] M. Žnidarič, Phys. Rev. B 88, 205135 (2013).
- [66] C. Karrasch, D. M. Kennes, and F. Heidrich-Meisner, Phys. Rev. B 91, 115130 (2015).
- [67] R. Steinigeweg, J. Herbrych, X. Zotos, and W. Brenig, Phys. Rev. Lett. **116**, 017202 (2016).
- [68] S. Nishimoto and M. Arikawa, Phys. Rev. B 79, 113106 (2009).
- [69] A. Metavitsiadis and W. Brenig, Phys. Rev. B 96, 041115(R) (2017).
- [70] M. Žnidarič, Phys. Rev. Lett. 110, 070602 (2013).
- [71] R. Steinigeweg, F. Heidrich-Meisner, J. Gemmer, K. Michielsen, and H. De Raedt, Phys. Rev. B 90, 094417 (2014).
- [72] C. Bartsch and J. Gemmer, Phys. Rev. Lett. 102, 110403 (2009).
- [73] P. Reimann, Phys. Rev. E 97, 062129 (2018).
- [74] K. Fabricius, U. Löw, and J. Stolze, Phys. Rev. B 55, 5833 (1997).
- [75] R. Steinigeweg and J. Gemmer, Phys. Rev. B 80, 184402 (2009).
- [76] For $\rho_l = S_{l,1}^z + S_{l,2}^z$ we have $\chi = 1/2$, while for $\rho_l = h_l$ we find $\chi = 3(2J_{\parallel}^2 + J_{\perp}^2)/16$.

- [77] Note that in the case of energy dynamics, we also find two smaller peaks at adjacent lattice sites $L/2 \pm 1$. This is caused by the fact that $[h_l, h_{l\pm 1}] \neq 0$.
- [78] R. Steinigeweg, H. Wichterich, and J. Gemmer, Europhys. Lett. (Europhys. Lett.) 88, 10004 (2009).
- [79] Y. Yan, F. Jiang, and H. Zhao, Eur. Phys. J. B 88, 11 (2015).
- [80] H. De Raedt and K. Michielsen, in *Handbook of Theoretical and Computational Nanotechnology* (American Scientific Publishers, Los Angeles, 2006).
- [81] V. V. Dobrovitski and H. A. De Raedt, Phys. Rev. E 67, 056702 (2003).

- [82] A. Weiße, G. Wellein, A. Alvermann, and H. Fehske, Rev. Mod. Phys. 78, 275 (2006).
- [83] V. K. Varma, A. Lerose, F. Pietracaprina, J. Goold, and A. Scardicchio, J. Stat. Mech. (2017) 053101.
- [84] J. Richter and R. Steinigeweg, Phys. Rev. E 99, 012114 (2019).
- [85] L. Knipschild and J. Gemmer, Phys. Rev. E 98, 062103 (2018).
- [86] L. Knipschild and J. Gemmer, Phys. Rev. A 99, 012118 (2019).
- [87] J. Herbrych, R. Steinigeweg, and P. Prelovšek, Phys. Rev. B 86, 115106 (2012).
- [88] I. Rousochatzakis, S. Kourtis, J. Knolle, R. Moessner, and N. B. Perkins, arXiv:1811.01671.